

\* **Integration by Parts**  

$$\int u dv = uv - \int v du$$

\* **Integral of Log**  

$$\int \ln x dx = x \ln x - x + C.$$

\* **Taylor Series**  
 If the function  $f$  is "smooth" at  $x = a$ , then it can be approximated by the  $n^{\text{th}}$  degree polynomial  

$$f(x) \approx f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \dots + \frac{f^{(n)}(a)}{n!}(x-a)^n.$$

\* **Maclaurin Series**  
 A Taylor Series about  $x = 0$  is called Maclaurin.  

$$e^x = 1 + x + \frac{x^2}{2} + \frac{x^3}{3!} + \dots$$

$$\cos(x) = 1 - \frac{x^2}{2} + \frac{x^4}{4!} - \dots$$

$$\sin(x) = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots$$

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots$$

$$\ln(x+1) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$$

\* **Lagrange Error Bound**  
 If  $P_n(x)$  is the  $n^{\text{th}}$  degree Taylor polynomial of  $f(x)$  about  $c$  and  $|f^{(n+1)}(t)| \leq M$  for all  $t$  between  $x$  and  $c$ , then  

$$|f(x) - P_n(x)| \leq \frac{M}{(n+1)!} |x-c|^{n+1}$$

\* **Alternating Series Error Bound**  
 If  $S_N = \sum_{k=1}^N (-1)^k a_k$  is the  $N^{\text{th}}$  partial sum of a convergent alternating series, then  

$$|S_\infty - S_N| \leq |a_{N+1}|$$

**Euler's Method**  
 If given that  $\frac{dy}{dx} = f(x,y)$  and that the solution passes through  $(x_0, y_0)$ ,  

$$y(x_0) = y_0$$
  

$$\vdots$$
  

$$y(x_n) = y(x_{n-1}) + f(x_{n-1}, y_{n-1}) \cdot \Delta x$$
  
 In other words:  

$$x_{\text{new}} = x_{\text{old}} + \Delta x$$
  

$$y_{\text{new}} = y_{\text{old}} + \left. \frac{dy}{dx} \right|_{(x_{\text{old}}, y_{\text{old}})} \cdot \Delta x$$

\* **Ratio Test**  
 The series  $\sum_{k=0}^{\infty} a_k$  converges if  

$$\lim_{k \rightarrow \infty} \left| \frac{a_{k+1}}{a_k} \right| < 1.$$
  
 If limit equals 1, you know nothing.

\* **Polar Curves**  
 For a polar curve  $r(\theta)$ , the **Area** inside a "leaf" is  

$$\int_{\theta_1}^{\theta_2} \frac{1}{2} [r(\theta)]^2 d\theta,$$
  
 where  $\theta_1$  and  $\theta_2$  are the "first" two times that  $r = 0$ .  
 The **slope** of  $r(\theta)$  at a given  $\theta$  is  

$$\frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta} = \frac{\frac{d}{d\theta}[r(\theta) \sin \theta]}{\frac{d}{d\theta}[r(\theta) \cos \theta]}$$

**L'Hopital's Rule**  
 If  $\frac{f(a)}{g(a)} = \frac{0}{0}$  or  $\frac{\infty}{\infty}$ ,  
 then  $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$ .

**VALUES OF TRIGONOMETRIC FUNCTIONS FOR COMMON ANGLES**

| $\theta$          | $\sin \theta$ | $\cos \theta$ | $\tan \theta$ |
|-------------------|---------------|---------------|---------------|
| $0^\circ$         | 0             | 1             | 0             |
| $\pi/6, 30^\circ$ | 1/2           | $\sqrt{3}/2$  | $\sqrt{3}/3$  |
| $37^\circ$        | 3/5           | 4/5           | 3/4           |
| $\pi/4, 45^\circ$ | $\sqrt{2}/2$  | $\sqrt{2}/2$  | 1             |
| $53^\circ$        | 4/5           | 3/5           | 4/3           |
| $\pi/3, 60^\circ$ | $\sqrt{3}/2$  | 1/2           | $\sqrt{3}$    |
| $\pi/2, 90^\circ$ | 1             | 0             | " $\infty$ "  |

\* means this is primarily for the BC exam

AP CALCULUS  
Stuff you **MUST** Know Cold

**Curve sketching and analysis**

$y = f(x)$  must be continuous at each:

critical point:  $\frac{dy}{dx} = 0$  or undefined.

local minimum : or endpoints

$\frac{dy}{dx}$  goes  $(-,0,+)$  or  $(-,und,+)$

or  $\frac{d^2y}{dx^2} > 0$ .

local maximum :

$\frac{dy}{dx}$  goes  $(+,0,-)$  or  $(+,und,-)$

or  $\frac{d^2y}{dx^2} < 0$ .

pt of inflection : concavity changes.

$\frac{d^2y}{dx^2}$  goes  $(+,0,-)$ ,  $(-,0,+)$ ,

$(+,und,-)$ , or  $(-,und,+)$

**Differentiation Rules**

Chain Rule

$$\frac{d}{dx} [f(u)] = f'(u) \frac{du}{dx}$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$$

Product Rule

$$\frac{d}{dx} (uv) = u \frac{dv}{dx} + \frac{du}{dx} v$$

Quotient Rule

$$\frac{d}{dx} \left( \frac{u}{v} \right) = \frac{\frac{du}{dx} v - u \frac{dv}{dx}}{v^2}$$

**Theorem of the Mean Value**  
i.e. Average Value

If the function  $f(x)$  is continuous on  $[a, b]$  and the first derivative exist on the interval  $(a, b)$ , then there exists a number  $x = c$  on  $(a, b)$  such that

$$f'(c) = \frac{\int_a^b f(x) dx}{(b-a)}$$

This value  $f(c)$  is the "average value" of the function on the interval  $[a, b]$ .

**Trapezoidal Rule**

$$\int_a^b f(x) dx = \frac{b-a}{2n} [f(x_0) + 2f(x_1) + \dots + 2f(x_{n-1}) + f(x_n)]$$

**Basic Derivatives**

$$\frac{d}{dx} (x^n) = nx^{n-1}$$

$$\frac{d}{dx} (\sin x) = \cos x$$

$$\frac{d}{dx} (\cos x) = -\sin x$$

$$\frac{d}{dx} (\tan x) = \sec^2 x$$

$$\frac{d}{dx} (\cot x) = -\csc^2 x$$

$$\frac{d}{dx} (\sec x) = \sec x \tan x$$

$$\frac{d}{dx} (\csc x) = -\csc x \cot x$$

$$\frac{d}{dx} (\ln x) = \frac{1}{x}$$

$$\frac{d}{dx} (e^x) = e^x$$

**"PLUS A CONSTANT"**

**The Fundamental Theorem of Calculus**

$$\int_a^b f(x) dx = F(b) - F(a)$$

where  $F'(x) = f(x)$ .

**Corollary to FTC**

$$\frac{d}{dx} \int_{a(x)}^{b(x)} f(t) dt = f(b(x)) b'(x) - f(a(x)) a'(x)$$

**Intermediate Value Theorem**

If the function  $f(x)$  is continuous on  $[a, b]$ , then for any number  $c$  between  $f(a)$  and  $f(b)$ , there exists a number  $d$  in the open interval  $(a, b)$  such that  $f(d) = c$ .

**Rolle's Theorem**

If the function  $f(x)$  is continuous on  $[a, b]$ , the first derivative exist on the interval  $(a, b)$ , and  $f(a) = f(b)$ ; then there exists a number  $x = c$  on  $(a, b)$  such that

$$f'(c) = 0.$$

**Mean Value Theorem**

If the function  $f(x)$  is continuous on  $[a, b]$ , and the first derivative exists on the interval  $(a, b)$ , then there exists a number  $x = c$  on  $(a, b)$  such that

$$f'(c) = \frac{f(b) - f(a)}{b - a}.$$

**Solids of Revolution and friends**

Disk Method

$$V = \pi \int_a^b [R(x)]^2 dx$$

Washer Method

$$V = \pi \int_a^b ([R(x)]^2 - [r(x)]^2) dx$$

Shell Method (no longer on AP)

$$V = 2\pi \int_a^b r(x) h(x) dx$$

ArcLength

$$L = \int_a^b \sqrt{1 + [f'(x)]^2} dx$$

Surface of revolution (No longer on AP)

$$S = 2\pi \int_a^b r(x) \sqrt{1 + [f'(x)]^2} dx$$

**More Derivatives**

$$\frac{d}{dx} (\sin^{-1} x) = \frac{1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx} (\cos^{-1} x) = \frac{-1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx} (\tan^{-1} x) = \frac{1}{1+x^2}$$

$$\frac{d}{dx} (\cot^{-1} x) = \frac{-1}{1+x^2}$$

$$\frac{d}{dx} (\sec^{-1} x) = \frac{1}{|x|\sqrt{x^2-1}}$$

$$\frac{d}{dx} (\csc^{-1} x) = \frac{-1}{|x|\sqrt{x^2-1}}$$

$$\frac{d}{dx} (a^x) = a^x \ln a$$

$$\frac{d}{dx} (\log_a x) = \frac{1}{x \ln a}$$

**Distance, velocity and acceleration**

velocity =  $\frac{d}{dt}$  (position).

acceleration =  $\frac{d}{dt}$  (velocity).

velocity vector =  $\left\langle \frac{dx}{dt}, \frac{dy}{dt} \right\rangle$ .

speed =  $|v| = \sqrt{(x')^2 + (y')^2}$ .

Distance =  $\int_{\text{initial time}}^{\text{final time}} |v| dt$

$$= \int_{t_0}^{t_f} \sqrt{(x')^2 + (y')^2} dt$$

average velocity =

$$\frac{\text{final position} - \text{initial position}}{\text{total time}}.$$